## This Page Is Inserted by IFW Operations and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**


Defective images within this document are accurate representations of the original documents submitted by the applicant.


Defects in the images may include (but are not limited to):


- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

## IMAGES ARE BEST AVAILABLE COPY.


As rescanning documents will not correct images, please do not report the images to the Image Problem Mailbox.







Cell attachment Heparanase T T 1E 1E 2E 2E kDa - + - + - + Cb Cb Cc



T = Trypsin

1E = 1mM EDTA


2E = 2mM EDTA

Cb = Control, purified heparanase from baculovirus, p60

Cc = Control, purified heparanase from CHO cells, p45


Fig. 2b

## Heparanase adherence to cells requires the presence of GAGs



Sample	Sample O.D sample		Δ O.D.	
B16-F1	0.1245	0.0422	0.0823	
B16-F1 transfected	0.1671	0.0303	0.1366	
B16-F1 adhered	0.4448	0.0321	0.4127	
Control b22	0.2258	0.1163	0.1095	

Fig. 4a



b22, b27, bmix = control, purified baculo heparanase (p60), different batches.
c = control, purified CHO heparanase (p45)
C = Control, substrate only

Fig. 4b




Fig. 4c

Sample #	Time of incubation with heparanase	Cell detachment from culture plate	O.D sample	Δ O.D.
1	0	Trypsin	0.1291	
2	15 min	Trypsin	0.1777	0.0486
3	60 min	Trypsin	0.318	0.1847
4	180 min	Trypsin	0.3633	0.2342
5	0	1mM EDTA	0.1370	
6	180 min	1mM EDTA	0.3754	0.2384

Fig. 5a




Fig. 5b




Fig. 6a





Fig. 6b



**Cb** = Control, from baculovirus, p60. **Cc** = Control, from CHO cells, p45.

Fig. 6c






Fig. 7








Fig. 8e

